



Health Visitor

Caroline Page & The Health Visiting Team are based at the Riverside Health Centre, James St. West, Bath BA1 2BT, and can be contacted on their direct line;

01225 474283

They have an answer phone and try to deal with messages daily. They are usually in the office between 8.30 and 9.30 to take phone calls.

They will offer advice and support to all families with babies and young children. Please contact them with any queries, or if you would like to arrange a home visit.

There is a weekly drop in child health clinic at Riverside Health Centre on Wednesdays between 11.00 am and 12.00 noon.

There are postnatal groups for new parents. Please ask us for details.

IMMUNISATION SCHEDULE

All children starting the immunisation programme at 2 months of age will follow the schedule below.

Immunisation appointments will be sent to you from the child health department. The children's immunisations are carried out by the practice nurse from 11.00 am - 12 noon on Wednesday mornings.

If you are unable to keep the appointment please contact the surgery to cancel. Once this has been done you will be recalled by the child health department.

IMPORTANT: Please remember to bring your baby's Child Health Surveillance (Red Book) with you to these appointments

UK 2011 Immunisation Schedule: taken from [http://www.patient.co.uk/doctor/Immunisation-Schedule-\(UK\).htm](http://www.patient.co.uk/doctor/Immunisation-Schedule-(UK).htm)

3 days	<ol style="list-style-type: none"> 1. BCG (if there has been tuberculosis in the family in the previous 6 months); 2. Hepatitis B vaccine if mother is HBsAg +ve. 	
2 months	<ol style="list-style-type: none"> 1. Diphtheria, tetanus, pertussis, polio and <i>Haemophilus influenzae</i> type b (DTaP/IPV/Hib). 2. Pneumococcal conjugate vaccine (PCV). 	<ol style="list-style-type: none"> 1. One injection (Pediacef®). 2. One injection (Prevenar13®).
3 months	<ol style="list-style-type: none"> 1. Diphtheria, tetanus, pertussis, polio and <i>H. influenzae</i> type b (DTaP/IPV/Hib). 2. Meningitis C (MenC). 	<ol style="list-style-type: none"> 1. One injection (Pediacef®). 2. One injection (NeisVac-C® or Meningitec®).
4 months	<ol style="list-style-type: none"> 1. Diphtheria, tetanus, pertussis, polio and <i>H. influenzae</i> type b (DTaP/IPV/Hib). 2. Pneumococcal conjugate vaccine (PCV). 3. Meningitis C (MenC). 	<ol style="list-style-type: none"> 1. One injection (Pediacef®). 2. One injection (Prevenar 13®). 3. One injection (NeisVac-C® or Meningitec®).
Between 12 and 13 months	<ol style="list-style-type: none"> 1. Measles, mumps and rubella (MMR). 2. Pneumococcal conjugate vaccine (PCV). 3. <i>H. influenzae</i> type b, meningitis C (Hib/MenC). 	<ol style="list-style-type: none"> 1. One injection (Priorix® or MMR II®). 2. One injection (Prevenar 13®) 3. One injection (Menitorix®).
3 years and 4 months to 5 years	<ol style="list-style-type: none"> 1. Diphtheria, tetanus, pertussis and polio (dTaP/IPV or DTaP/IPV). 2. Measles, mumps and rubella (MMR). 	<ol style="list-style-type: none"> 1. One injection (Repevax® or Infanrix-IPV®). 2. One injection (Priorix® or MMR II®).
Girls aged 12 to 13 years	<ul style="list-style-type: none"> • Cervical cancer caused by human papillomavirus (HPV) types 16 and 18. 	<ul style="list-style-type: none"> • HPV (Cervarix®) Three injections: (the second 1-2 months and third 6 months after first injection).
13-18 years	<ul style="list-style-type: none"> • Tetanus, diphtheria and polio (Td/IPV). 	<ul style="list-style-type: none"> • One injection (Revaxis®).